

# Criteria for subgeometric ergodicity of strong Markov processes

Gersende FORT

CNRS/ENST, France

Joint works with :

- ▶ Gareth ROBERTS (Lancaster University, UK).
- ▶ Randal DOUC (Ecole Polytechnique, France) & Arnaud GUILLIN (Université Paris IX, France).

# Criteria for subgeometric ergodicity of strong Markov processes

Gersende FORT

CNRS/ENST, France

Joint works with :

- ▶ Gareth ROBERTS (Lancaster University, UK).
- ▶ Randal DOUC (Ecole Polytechnique, France) & Arnaud GUILLIN (Université Paris IX, France).

Let

- ▶  $\{\Phi_t\}_{t \geq 0}$  be a time-continuous **strong** Markov process on  $X$  ( $X = \mathbb{R}^d$ ), with **cadlag** paths.
- ▶  $\{P^t\}_{t \geq 0}$  be the Markov semigroup :  $\mathbb{P}_x(\Phi_t \in A) = P^t(x, A)$ .

# Objective

Conditions that imply

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| = 0 \quad \forall x \in X, \quad (1)$$

$\pi$  : (the) invariant probability measure

$f$  : positive function,

Ex.  $f = 1$  [Total variation norm],  $f(x) \sim |x|^p, \dots$

$r$  : positive non-decreasing rate function

Subgeometric rate function:

Ex.  $r(t) \sim t^\tau$   $r(t) \sim \{\log t\}^\tau$   $r(t) \sim \exp(ct^\gamma)$   $0 < \gamma < 1$ .

$$0 < \liminf_t r(t)/\tilde{r}(t) \leq \limsup_t r(t)/\tilde{r}(t) < \infty, \quad \lim_t \frac{\log \tilde{r}(t)}{t} \downarrow 0.$$

# Objective

Conditions that imply

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| = 0 \quad \forall x \in X, \quad (1)$$

$\pi$  : (the) invariant probability measure

$f$  : positive function,

Ex.  $f = 1$  [Total variation norm],  $f(x) \sim |x|^p, \dots$

$r$  : positive non-decreasing rate function

Subgeometric rate function:

Ex.  $r(t) \sim t^\tau$   $r(t) \sim \{\log t\}^\tau$   $r(t) \sim \exp(ct^\gamma)$   $0 < \gamma < 1$ .

$$0 < \liminf_t r(t)/\tilde{r}(t) \leq \limsup_t r(t)/\tilde{r}(t) < \infty, \quad \lim_t \frac{\log \tilde{r}(t)}{t} \downarrow 0.$$

► Answer :

(1)  $\Leftrightarrow$  Return-time to a petite set  $\Leftrightarrow$  Drift inequalities



Regular set and regular measures

Skeleton chain, resolvent

Moderate deviation principle

# Definitions

Delayed return-time to  $A$ :

$$\tau_A(\delta) = \inf\{t \geq \delta, \Phi_t \in A\}, \quad \tau_A(0) = \tau_A.$$

# Definitions

Delayed return-time to  $A$ :

$$\tau_A(\delta) = \inf\{t \geq \delta, \Phi_t \in A\}, \quad \tau_A(0) = \tau_A.$$

Phi-irreducibility: there exists a measure  $\phi$

$$\phi(A) > 0 \implies \forall x, \quad \mathbb{P}_x(\tau_A < \infty) > 0$$

# Definitions

Delayed return-time to  $A$ :

$$\tau_A(\delta) = \inf\{t \geq \delta, \Phi_t \in A\}, \quad \tau_A(0) = \tau_A.$$

Phi-irreducibility: there exists a measure  $\phi$

$$\phi(A) > 0 \implies \forall x, \quad \mathbb{P}_x(\tau_A < \infty) > 0$$

Maximal irred. measure: irred. measure  $\psi$  s.t.  $\phi \prec \psi$ .

Accessible set: a set s.t.  $\psi(A) > 0$ .

# Definitions

Delayed return-time to  $A$ :

$$\tau_A(\delta) = \inf\{t \geq \delta, \Phi_t \in A\}, \quad \tau_A(0) = \tau_A.$$

Phi-irreducibility: there exists a measure  $\phi$

$$\phi(A) > 0 \implies \forall x, \quad \mathbb{P}_x(\tau_A < \infty) > 0$$

Maximal irred. measure: irred. measure  $\psi$  s.t.  $\phi \prec \psi$ .

Accessible set: a set s.t.  $\psi(A) > 0$ .

Petite set  $C$ : there exist a probability measure  $a$  on  $[0, \infty)$  and  $\nu_a$  measure on  $X$

$$\int_0^\infty P^t(x, \cdot) a(dt) \geq \nu_a(\cdot), \quad \forall x \in C.$$



# Definitions

Delayed return-time to  $A$ :

$$\tau_A(\delta) = \inf\{t \geq \delta, \Phi_t \in A\}, \quad \tau_A(0) = \tau_A.$$

Phi-irreducibility: there exists a measure  $\phi$

$$\phi(A) > 0 \implies \forall x, \quad \mathbb{P}_x(\tau_A < \infty) > 0$$

Maximal irred. measure: irred. measure  $\psi$  s.t.  $\phi \prec \psi$ .

Accessible set: a set s.t.  $\psi(A) > 0$ .

Petite set  $C$ : there exist a probability measure  $a$  on  $[0, \infty)$  and  $\nu_a$  measure on  $X$

$$\int_0^\infty P^t(x, \cdot) a(dt) \geq \nu_a(\cdot), \quad \forall x \in C.$$

Existence: A phi-irreducible process possesses an accessible closed petite set.

# Definitions

Delayed return-time to  $A$ :

$$\tau_A(\delta) = \inf\{t \geq \delta, \Phi_t \in A\}, \quad \tau_A(0) = \tau_A.$$

Phi-irreducibility: there exists a measure  $\phi$

$$\phi(A) > 0 \implies \forall x, \quad \mathbb{P}_x(\tau_A < \infty) > 0$$

Maximal irred. measure: irred. measure  $\psi$  s.t.  $\phi \prec \psi$ .

Accessible set: a set s.t.  $\psi(A) > 0$ .

Petite set  $C$ : there exist a probability measure  $a$  on  $[0, \infty)$  and  $\nu_a$  measure on  $X$

$$\int_0^\infty P^t(x, \cdot) a(dt) \geq \nu_a(\cdot), \quad \forall x \in C.$$

Existence: A phi-irreducible process possesses an accessible closed petite set.

Aperiodicity: there exist an accessible petite set  $C$  and  $t_0 > 0$  such that

$$\inf_{x \in C} P^t(x, C) > 0, \quad \forall t \geq t_0.$$

# Discrete-time / Continuous-time

There exist a rate function  $r$  and a function  $f$  such that

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| = 0 \quad \forall x \in X,$$

Discrete-time case :

- phi-irreducible, aperiodic,
  - $C$  petite set such that  $\sup_{x \in C} \mathbb{E}_x \left[ \sum_{k=0}^{\tau_C - 1} \tilde{r}(k) f(\Phi_k) \right] < \infty$ ,
- when
- $r(t) = \tilde{r}(t) = 1$ ,
  - $r(t) \sim \kappa^t$ ,  $\kappa > 1$ ,  $r \neq \tilde{r}$  MEYN & TWEEDIE, 1993
  - $r = \tilde{r}$  subgeom. JARNER ROBERTS 2002, FORT MOULINES 2003, DOUC ET AL. 2004

# Discrete-time / Continuous-time

There exist a rate function  $r$  and a function  $f$  such that

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| = 0 \quad \forall x \in X,$$

Discrete-time case :

- $\phi$ -irreducible, aperiodic,
  - $C$  petite set such that  $\sup_{x \in C} \mathbb{E}_x \left[ \sum_{k=0}^{\tau_C - 1} \tilde{r}(k) f(\Phi_k) \right] < \infty$ ,
- when
- $r(t) = \tilde{r}(t) = 1$ ,
  - $r(t) \sim \kappa^t$ ,  $\kappa > 1$ ,  $r \neq \tilde{r}$  MEYN & TWEEDIE, 1993
  - $r = \tilde{r}$  subgeom. JARNER ROBERTS 2002, FORT MOULINES 2003, DOUC ET AL. 2004

Continuous-time case :

- $\phi$ -irreducible, aperiodic,
  - $C$  closed petite set s.t.  $\sup_{x \in C} \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} \tilde{r}(s) f(\Phi_s) ds \right] < \infty$ ,
- when
- $r(t) = \tilde{r}(t) = 1$  MEYN & TWEEDIE 1993
  - $r(t) \sim \kappa^t$ ,  $\kappa > 1$   $r \neq \tilde{r}$  MEYN TWEEDIE 1993, DOWN ET AL. 1995
  - $\hookrightarrow$  subgeometric rate function ?

# Res. 1

**Theorem (subgeometric  $f$ -ergodicity)** FORT & ROBERTS, 2005

- If**
- The process is Harris-recurrent with invariant prob. distribution  $\pi$ .
  - Some skeleton chain  $P^m$  is phi-irreducible.
  - There exist a closed petite set  $C$ ,  $\delta > 0$ , a function  $f_* \geq 1$  and a subgeometric rate function  $r_*$  s.t.

$$\sup_{x \in C} \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right] < \infty, \quad \sup_{x \in C} \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right] < \infty.$$

- There exists  $c$  such that  $\sup_{t \leq m} P^t f_* \leq c f_*$ .

# Res. 1

**Theorem (subgeometric  $f$ -ergodicity)** FORT & ROBERTS, 2005

- If**
- The process is Harris-recurrent with invariant prob. distribution  $\pi$ .
  - Some skeleton chain  $P^m$  is  $\phi$ -irreducible.
  - There exist a closed petite set  $C$ ,  $\delta > 0$ , a function  $f_* \geq 1$  and a subgeometric rate function  $r_*$  s.t.

$$\sup_{x \in C} \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right] < \infty, \quad \sup_{x \in C} \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right] < \infty.$$

- There exists  $c$  such that  $\sup_{t \leq m} P^t f_* \leq c f_*$ .

- Then**
- $\pi(f_*) < \infty$ .
  - $\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| = 0, \quad x \in \mathcal{S}_*$ ,

$$\begin{array}{lll} r(t) = r_*(t) & r(t) = 1 & r(t) = \Psi_1(r_*(t)) \\ f(t) = 1 & f(t) = f_*(t) & f(t) = \Psi_2(f_*(t)) \end{array}, \quad \Psi_1(x)\Psi_2(y) \leq x + y$$

- and similar conclusions

$$U_{r_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right], \quad U_{f_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right].$$

► If  $\mu$  is a probability measure such that  $\mu(U_{r_*} + U_{f_*}) < \infty$ ,

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} \left| \int \mu(dx) P^t g(x) - \pi(g) \right| = 0.$$

$$U_{r_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right], \quad U_{f_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right].$$

- If  $\mu$  is a probability measure such that  $\mu(U_{r_*} + U_{f_*}) < \infty$ ,

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} \left| \int \mu(dx) P^t g(x) - \pi(g) \right| = 0.$$

- For all  $x \in X$ ,

$$r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| \leq c \{U_{r_*}(x) + U_{f_*}(x)\}.$$



$$U_{r_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right], \quad U_{f_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right].$$

- If  $\mu$  is a probability measure such that  $\mu(U_{r_*} + U_{f_*}) < \infty$ ,

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} \left| \int \mu(dx) P^t g(x) - \pi(g) \right| = 0.$$

- For all  $x \in X$ ,

$$r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| \leq c \{U_{r_*}(x) + U_{f_*}(x)\}.$$

- For all  $x, y \in X$ ,

$$\int_0^\infty r(t) \sup_{|g| \leq f} |P^t g(x) - P^t g(y)| dt \leq c \{U_{r_*}(x) + U_{f_*}(x) + U_{r_*}(y) + U_{f_*}(y)\}.$$

$$U_{r_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right], \quad U_{f_*}(x) = \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right].$$

- If  $\mu$  is a probability measure such that  $\mu(U_{r_*} + U_{f_*}) < \infty$ ,

$$\lim_{t \rightarrow \infty} r(t) \sup_{|g| \leq f} \left| \int \mu(dx) P^t g(x) - \pi(g) \right| = 0.$$

- For all  $x \in X$ ,

$$r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| \leq c \{U_{r_*}(x) + U_{f_*}(x)\}.$$

- For all  $x, y \in X$ ,

$$\int_0^\infty r(t) \sup_{|g| \leq f} |P^t g(x) - P^t g(y)| dt \leq c \{U_{r_*}(x) + U_{f_*}(x) + U_{r_*}(y) + U_{f_*}(y)\}.$$

- If  $\partial r$  is a subgeometric rate function,

$$\int_0^\infty \partial r(t) \sup_{|g| \leq f} |P^t g(x) - \pi(g)| dt \leq C \{U_{r_*}(x) + U_{f_*}(x)\}.$$

## Res. 2 : Sufficient Conditions for delayed return-time

$$\text{Find } \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right] \leq V_{r_*}(x), \quad \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right] \leq V_{f_*}(x).$$

**Theorem (Answer 1)** DOUC, FORT & GULLIN, 2006

**If** For all  $x \in X$ ,

$s \mapsto V(\Phi_s) - V(X_0) - \int_0^s \{-\phi \circ V(\Phi_u) + b \mathbb{1}_C(u)\} du$  is a  $\mathbb{P}_x$ -supermartingale

$C$  closed set,  $V \geq 1$  cadlag,  $\phi > 0 \uparrow$  differentiable concave function.

## Res. 2 : Sufficient Conditions for delayed return-time

$$\text{Find } \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_*(s) ds \right] \leq V_{r_*}(x), \quad \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} f_*(\Phi_s) ds \right] \leq V_{f_*}(x).$$

**Theorem (Answer 1)** DOUC, FORT & GUILIN, 2006

**If** For all  $x \in X$ ,

$s \mapsto V(\Phi_s) - V(X_0) - \int_0^s \{-\phi \circ V(\Phi_u) + b \mathbb{1}_C(u)\} du$  is a  $\mathbb{P}_x$ -supermartingale

$C$  closed set,  $V \geq 1$  cadlag,  $\phi > 0 \uparrow$  differentiable concave function.

**Then** for all  $x \in X$ ,  $\delta > 0$ ,

$$\mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} r_\phi(s) ds \right] \leq \tilde{c}_\delta V(x), \quad \mathbb{E}_x \left[ \int_0^{\tau_C(\delta)} \phi \circ V(\Phi_s) ds \right] \leq c_\delta V(x),$$

$$\text{where } r_\phi(s) = \phi \circ H_\phi^{-1}(s), \quad H_\phi(s) = \int_1^s \frac{du}{\phi(u)}.$$

**Rmk :**  $r_\phi$  subgeometric rate function if  $\lim_{t \rightarrow \infty} \phi'(t) = 0$ .

## Res. 2 : Sufficient conditions for delayed return-time

Let  $\mathcal{A}$  be the extended generator with domain  $\mathcal{D}(\mathcal{A}) : V \in \mathcal{D}(\mathcal{A})$  iff

$$s \mapsto V(\Phi_s) - V(\Phi_0) - \int_0^s \mathcal{A}V(\Phi_u) du \quad \mathbb{P}_x\text{-local martingale.}$$

**Theorem (Answer 2)** FORT & ROBERTS 2005; DOUC, FORT & GUILLIN, 2006

**If** for all  $x \in X$ ,

$$\mathcal{A}V(x) \leq -\phi \circ V(x) + b\mathbb{1}_C(x),$$

$C$  closed set,  $V \geq 1$  cadlag in  $\mathcal{D}(\mathcal{A})$ ,  $\phi > 0$  increasing differentiable concave function.

**Then** the previous theorem applies :

$$r_*(s) = r_\phi(s), \quad f_*(x) = \phi \circ V(x).$$

## Res. 2 : Sufficient conditions for delayed return-time

Let  $\mathcal{A}$  be the extended generator with domain  $\mathcal{D}(\mathcal{A}) : V \in \mathcal{D}(\mathcal{A})$  iff

$$s \mapsto V(\Phi_s) - V(\Phi_0) - \int_0^s \mathcal{A}V(\Phi_u) du \quad \mathbb{P}_x\text{-local martingale.}$$

**Theorem (Answer 2)** FORT & ROBERTS 2005; DOUC, FORT & GUILLIN, 2006

If for all  $x \in X$ ,

$$\mathcal{A}V(x) \leq -\phi \circ V(x) + b\mathbb{1}_C(x),$$

$C$  closed set,  $V \geq 1$  cadlag in  $\mathcal{D}(\mathcal{A})$ ,  $\phi > 0$  increasing differentiable concave function.

Then the previous theorem applies :

$$r_*(s) = r_\phi(s), \quad f_*(x) = \phi \circ V(x).$$

**Example**

DOUC ET AL. 2004

- $\phi(t) \sim t^{1-\alpha} \longrightarrow r_\phi(t) \sim t^{1/\alpha-1} \quad 0 < \alpha < 1.$
- $\phi(t) \sim (1 + \log t)^\alpha \longrightarrow r_\phi(t) \sim \log^\alpha(t) \quad \alpha > 0.$
- $\phi(t) \sim t/(\log t)^\alpha \longrightarrow r_\phi(t) \sim t^{-\alpha/(1+\alpha)} \exp(ct^{1/(1+\alpha)}) \quad \alpha > 0.$

## Drift inequalities : continuous-time / discrete-time

	Continuous-time case	Discrete-time case
Non explosivity	$\mathcal{A}V \leq cV$	
Recurrence	$\mathcal{A}V \leq c\mathbb{1}_C$	$PV - V \leq b\mathbb{1}_C$
$f$ -ergodicity	$\mathcal{A}V \leq -f + b\mathbb{1}_C$	$PV - V \leq -f + b\mathbb{1}_C$
Geometric ergodicity	$\mathcal{A}V \leq -cV + b\mathbb{1}_C$	$PV - V \leq -cV + b\mathbb{1}_C \quad 0 < c < 1$
Polynomial ergodicity		$PV - V \leq -cV^{1-\alpha} + b\mathbb{1}_C$
Subgeometric ergodicity		$PV - V \leq -\phi \circ V + b\mathbb{1}_C$

## Drift inequalities : continuous-time / discrete-time

	Continuous-time case	Discrete-time case
Non explosivity	$\mathcal{A}V \leq cV$	
Recurrence	$\mathcal{A}V \leq c\mathbb{1}_C$	$PV - V \leq b\mathbb{1}_C$
$f$ -ergodicity	$\mathcal{A}V \leq -f + b\mathbb{1}_C$	$PV - V \leq -f + b\mathbb{1}_C$
Geometric ergodicity	$\mathcal{A}V \leq -cV + b\mathbb{1}_C$	$PV - V \leq -cV + b\mathbb{1}_C \quad 0 < c < 1$
Polynomial ergodicity	$\mathcal{A}V \leq -cV^{1-\alpha} + b\mathbb{1}_C$	$PV - V \leq -cV^{1-\alpha} + b\mathbb{1}_C$
Subgeometric ergodicity	$\mathcal{A}V \leq -\phi \circ V + b\mathbb{1}_C$	$PV - V \leq -\phi \circ V + b\mathbb{1}_C$



## Example: Elliptic diffusions on $\mathbb{R}^n$ (a)

Consider

$$\Phi_t = \Phi_0 + \int_0^t b(\Phi_s) ds + \int_0^t \sigma(\Phi_s) dB_s,$$

### Conditions on $b, \sigma$

- $\sigma$  bounded;  $b, \sigma$  locally lipschitz.
- $a(x)$  non singular.
- There exist  $0 < p < 1, \quad M, r > 0$  s.t.

$$\langle b(x), x \rangle \leq -r|x|^{1-p} \quad |x| \geq M.$$

## Example: Elliptic diffusions on $\mathbb{R}^n$ (a)

Consider

$$\Phi_t = \Phi_0 + \int_0^t b(\Phi_s) ds + \int_0^t \sigma(\Phi_s) dB_s,$$

### Conditions on $b, \sigma$

- $\sigma$  bounded;  $b, \sigma$  locally lipschitz.
- $a(x)$  non singular.
- There exist  $0 < p < 1$ ,  $M, r > 0$  s.t.

$$\langle b(x), x \rangle \leq -r|x|^{1-p} \quad |x| \geq M.$$

### Drift inequalities

Itô's formula yields

$$\mathcal{A}V(x) = \langle b(x), \nabla V(x) \rangle + \frac{1}{2} \text{Tr} (\nabla^2 V(x) a(x)),$$

Ex.  $V(x) = \exp(\iota|x|^m)$ .

## Example: Elliptic diffusions on $\mathbb{R}^n$ (b)

Set  $\lambda_+ := \sup_{x \neq 0} |x|^2 \langle a(x)x, x \rangle.$

**Theorem (Ergodicity of the diffusion)**

$\pi$ -integrable  $\exists \pi$  and for all  $c > 0$  s.t.  $r - 0.5c\lambda_+(1-p) > 0$

$$\int \pi(dx) \exp(c|x|^{1-p}) < \infty.$$

## Example: Elliptic diffusions on $\mathbb{R}^n$ (b)

Set  $\lambda_+ := \sup_{x \neq 0} |x|^2 \langle a(x)x, x \rangle$ .

### Theorem (Ergodicity of the diffusion)

$\pi$ -integrable  $\exists \pi$  and for all  $c > 0$  s.t.  $r - 0.5c\lambda_+(1-p) > 0$

$$\int \pi(dx) \exp(c|x|^{1-p}) < \infty.$$

Subgeometric ergodicity for all  $c > 0$  s.t.  $r - 0.5c\lambda_+(1-p) > 0$

$$r_*(t) \sim t^{-2p/(1+p)} \exp(\{\tilde{c}t\}^{(1-p)/(1+p)}), \quad f_*(x) \sim |x|^{-2p} \exp(c|x|^{1-p}),$$

and  $\tilde{c} = c^{(1+p)/(1-p)(1+p)}\{r - 0.5c\lambda_+(1-p)\}$ .

## Example: Langevin Tempered diffusions on $\mathbb{R}^n$ (a)

Let  $\pi$  be a probability measure. Consider

$$\Phi_t = \Phi_0 + \int_0^t b(\Phi_s) ds + \int_0^t \sigma(\Phi_s) dB_s,$$

where

$$b(x) = \frac{1}{2} \sum_{j=1}^n a_{i,j}(x) \partial_{x_j} \log \pi(x) + \frac{1}{2} \sum_{j=1}^n \partial_{x_j} a_{i,j}(x) \quad a(x) = \sigma(x)\sigma'(x).$$

## Example: Langevin Tempered diffusions on $\mathbb{R}^n$ (a)

Let  $\pi$  be a probability measure. Consider

$$\Phi_t = \Phi_0 + \int_0^t b(\Phi_s) ds + \int_0^t \sigma(\Phi_s) dB_s,$$

where

$$b(x) = \frac{1}{2} \sum_{j=1}^n a_{i,j}(x) \partial_{x_j} \log \pi(x) + \frac{1}{2} \sum_{j=1}^n \partial_{x_j} a_{i,j}(x) \quad a(x) = \sigma(x)\sigma'(x).$$

**Problem** :  $\pi$  is heavy tailed

• if  $\sigma(x) = c|x|^{-d}$ , can not be geometrically ergodic ROBERTS TWEEDIE

• What happens if  $\sigma(x) = \pi^{-d}(x)$ ,  $d > 0$  ?

## Example: Langevin Tempered diffusions on $\mathbb{R}^n$ (b)

### Conditions

#### On $\pi$

- $\pi$  is positive and  $C^2$  on  $\mathbb{R}^n$ .
- $\pi$  is polynomially decreasing in the tails :  $\exists 0 < \beta < 1/n$

$$0 < \liminf_{|x| \rightarrow \infty} \frac{|\nabla \log \pi(x)|}{\pi^\beta(x)} \leq \limsup_{|x| \rightarrow \infty} \frac{|\nabla \log \pi(x)|}{\pi^\beta(x)} < \infty,$$

$$2\beta - 1 < \gamma := \liminf_{|x| \rightarrow \infty} \frac{\text{Tr}(\nabla^2 \log \pi(x))}{|\nabla \log \pi(x)|^2} \leq \limsup_{|x| \rightarrow \infty} \frac{\text{Tr}(\nabla^2 \log \pi(x))}{|\nabla \log \pi(x)|^2} < \infty,$$

## Example: Langevin Tempered diffusions on $\mathbb{R}^n$ (b)

### Conditions

#### On $\pi$

- $\pi$  is positive and  $C^2$  on  $\mathbb{R}^n$ .
- $\pi$  is polynomially decreasing in the tails :  $\exists 0 < \beta < 1/n$

$$0 < \liminf_{|x| \rightarrow \infty} \frac{|\nabla \log \pi(x)|}{\pi^\beta(x)} \leq \limsup_{|x| \rightarrow \infty} \frac{|\nabla \log \pi(x)|}{\pi^\beta(x)} < \infty,$$

$$2\beta - 1 < \gamma := \liminf_{|x| \rightarrow \infty} \frac{\text{Tr}(\nabla^2 \log \pi(x))}{|\nabla \log \pi(x)|^2} \leq \limsup_{|x| \rightarrow \infty} \frac{\text{Tr}(\nabla^2 \log \pi(x))}{|\nabla \log \pi(x)|^2} < \infty,$$

#### On $d$

- $d \in \mathcal{D}_n$  s.t. the process is non-explosive.

**Example:**  $\pi(x) \sim |x|^{-1/\beta}$

$$\mathcal{D}_1 = [0; (1+\beta)/2] \quad \mathcal{D}_n = [0; (1+\beta(2-n))/2]; \quad \gamma = \beta(2-n) > 2\beta - 1.$$



## Example: Langevin Tempered diffusions on $\mathbb{R}^n$ (c)

Itô's formula yields

$$\mathcal{A}V(x) = \langle b(x), \nabla V(x) \rangle + \frac{1}{2} \text{Tr} (\nabla^2 V(x) a(x)).$$

**Theorem (Ergodicity of the Langevin tempered diffusion)** If  $d \in \mathcal{D}_n$  and

$0 \leq d < \beta$ : Fails to be geometrically ergodic.

Is polynomially ergodic, for all  $0 \leq \kappa < 1 + \gamma - 2\beta$

$$\lim_t (1+t)^\tau \sup_{|g| \leq 1 + \pi^{-\kappa}} |P^t g(x) - \pi(g)| = 0, \quad 0 \leq \tau < \frac{1 + \gamma - 2\beta - \kappa}{2(\beta - d)}.$$

## Example: Langevin Tempered diffusions on $\mathbb{R}^n$ (c)

Itô's formula yields

$$\mathcal{A}V(x) = \langle b(x), \nabla V(x) \rangle + \frac{1}{2} \text{Tr} (\nabla^2 V(x) a(x)).$$

**Theorem (Ergodicity of the Langevin tempered diffusion)** If  $d \in \mathcal{D}_n$  and

$0 \leq d < \beta$ : Fails to be geometrically ergodic.

Is polynomially ergodic, for all  $0 \leq \kappa < 1 + \gamma - 2\beta$

$$\lim_t (1+t)^\tau \sup_{|g| \leq 1 + \pi^{-\kappa}} |P^t g(x) - \pi(g)| = 0, \quad 0 \leq \tau < \frac{1 + \gamma - 2\beta - \kappa}{2(\beta - d)}.$$

$0 < \beta \leq d < (1 + \gamma)/2$ : Is geometrically ergodic, for all

$0 < \kappa < 1 + \gamma - 2d$ ,

$$\exists \tau > 1 \quad \lim_t \tau^t \sup_{|g| \leq 1 + \pi^{-\kappa}} |P^t g(x) - \pi(g)| = 0.$$

$0 < \beta < d$ : Is uniformly ergodic,

$$\exists \tau > 1 \text{ and } c \quad \forall x \in X, \quad \lim_t \tau^t \sup_{|g| \leq 1} |P^t g(x) - \pi(g)| \leq c.$$

# Conclusion

## Other results

When

$$s \mapsto V(\Phi_s) - V(\Phi_0) - \int_0^s \{-\phi \circ V(\Phi_u) + b\mathbb{1}_C(u)\} du \quad \mathbb{P}_x\text{-supermartingale,}$$
$$\sup_C V < \infty$$

where  $C$  closed petite set,  $V \geq 1$  cadlag,  $\phi \uparrow$  differentiable concave.

### Petite set

- the level sets  $\{V \leq n\}$  are petite.

### $(f, r)$ -regularity

- the level sets  $\{V \leq n\}$  are  $(f, r)$ -regular.
- there exists a full set, union of  $(f, r)$ -regular sets.

### Skeleton

- Subgeometric moment of the return-time to a small set, for **any** skeleton.

### Resolvent

- Subgeometric condition for **any** resolvent kernel.

### Moderate deviation principle ...

# Conclusion

Details can be found in:

- G. Fort & G.O. Roberts. Subgeometric ergodicity of strong Markov processes. *Ann. Appl. Probab.* 15(2):1565-1589, 2005.
- R. Douc, G. Fort & A. Guillin. Subgeometric rates of convergence of  $f$ -ergodic strong Markov processes. *ArXiv math.ST/0605791*, 2006.